

Sample size: 1122

Adult	HSR 21: Jefferson			Colorado		
	2008 Estimates (Weighted Percent)	Lower CI	Upper CI	2008 Estimates (Weighted Percent)	Lower CI	Upper CI
Smoking status						
Current smoking	16.7	13.0	20.4	18.1	16.9	19.4
by sex						
men	18.1	12.2	24.0	20.6	18.6	22.7
women	15.3	10.8	19.9	15.7	14.3	17.0
by age						
18-24	23.9	8.1	39.7	25.1	19.8	30.3
25-64	17.2	13.1	21.3	18.6	17.2	20.0
65+	7.3	4.1	10.5	8.8	7.2	10.3
by race/ethnicity						
white (non-Hispanic)	17.8	13.9	21.6	16.8	15.5	18.1
Latino/Hispanic	15.0	0.5	29.5	26.9	20.9	32.9
African-American/black	–	–	–	13.0	8.9	17.1
Asian American/Native Hawaiian/other Pacific Islander	–	–	–	20.2	14.6	25.7
American Indian/Alaska Native	–	–	–	42.6	31.5	53.8
all other	–	–	–	19.4	13.0	25.8
by socio-economic disadvantage*						
yes	35.3	24.7	45.8	29.2	26.6	31.8
no	9.9	7.1	12.6	11.4	10.1	12.7
Never-smoker	54.6	49.8	59.3	56.6	55.2	58.1
Former smoker	28.7	24.5	32.9	25.2	24.1	26.4
Daily smokers (of current)	72.3	59.2	85.4	70.9	66.9	74.9
Among quit attempters in region:						
used NRT	25.8	14.6	37.0	19.9	16.9	22.9
used Chantix	10.8	3.3	18.2	6.4	4.8	8.1
heard of QuitLine	91.5	81.3	101.7	83.7	80.7	86.6
ever called QuitLine	17.1	9.1	25.2	12.8	10.6	15.0
called QuitLine in past year	12.4	5.4	19.5	7.4	5.7	9.0
Among smokers in the region during the past year:						
made a quit attempt (of smokers during past year)	63.8	53.0	74.6	65.4	61.8	69.0
were advised to quit by provider	63.2	49.2	77.2	61.3	56.9	65.7
were referred to quitting help by provider, of those who were advised	38.2	25.8	50.6	41.2	36.3	46.1
Non-cigarette tobacco use						
current spit tobacco user	1.8	0.8	2.9	3.1	2.5	3.6
men	3.8	1.6	5.9	5.8	4.8	6.9
women	0.0	0.0	0.0	0.3	0.1	0.5
Secondhand smoke						
have rule that smoking is not allowed in the home	87.0	83.7	90.3	85.4	84.3	86.5
in households with smokers	73.7	65.9	81.5	63.7	60.7	66.7
in households without smokers	92.7	88.9	96.5	94.3	93.6	95.0
have rule that smoking is not allowed in the car	80.9	76.8	84.9	79.9	78.5	81.2
in households with smokers	53.3	40.3	66.2	47.4	43.8	51.0
in households without smokers	91.0	87.6	94.4	91.1	90.0	92.2
someone smoked at work in past 30 days	3.4	1.8	5.1	5.2	4.2	6.3
someone smoked near you in past 6 months (outside of work or home)	42.8	38.0	47.6	39.6	38.2	41.1

*Socio-economic disadvantage is defined as: <HS education, no insurance, <200% FPL or being disabled
 –Cells with a sample size smaller than 25 may yield unreliable results and have therefore been suppressed

Sample size: 726		HSR 21: Jefferson			Colorado		
Youth	2008 Estimates (Weighted Percent)	Lower CI	Upper CI	2008 Estimates (Weighted Percent)	Lower CI	Upper CI	
Smoking status among high schoolers							
Ever-smoker	38.7	33.2	44.2	36.8	35.2	38.3	
Current smoker	14.5	7.7	21.3	11.9	10.7	13.2	
by sex							
male	18.4	9.7	27.2	13.1	11.5	14.7	
female	11.2	5.3	17.0	10.9	9.6	12.1	
by ethnicity							
white (non-Hispanic)	15.1	7.1	23.1	12.3	10.6	13.9	
Latino/Hispanic	12.2	6.2	18.2	12.6	11.2	14.1	
African-American/black	–	–	–	6.0	2.9	9.0	
Asian American/Native							
Hawaiian/other Pacific Islander	7.6	-1.7	16.9	7.6	4.7	10.5	
American Indian/Alaska Native	–	–	–	24.4	14.8	34.1	
all other	26.0	13.8	38.3	12.6	9.1	16.2	
Among smokers in high school during the past year:							
frequent smoker (smoked 20-30 days in past 30)	6.2	4.1	8.3	5.3	4.7	5.8	
made a quit attempt in past year	54.0	46.7	61.3	51.7	48.3	55.0	
Of never smokers in high school:							
susceptible to smoking in next year	14.9	11.1	18.7	19.1	17.4	20.8	
Non-cigarette tobacco use among high schoolers							
current chew user	9.5	5.8	13.3	7.4	6.4	8.4	
current cigar user	17.3	9.5	25.2	15.0	13.7	16.3	
ever other tobacco user	24.9	17.4	32.3	19.7	18.1	21.2	
Secondhand smoke (SHS) among high schoolers							
exposed to SHS in room, past 7 days	39.7	33.5	46.0	36.0	34.0	38.0	
exposed to SHS in car, past 7 days	35.4	28.1	42.7	33.1	31.1	35.2	
thinks SHS is harmful	61.2	56.9	65.5	62.4	59.8	65.0	

–Cells with a sample size smaller than 25 may yield unreliable results and have therefore been suppressed