



## Scentless chamomile

*(Tripleurosperum perforatum)*



Scentless chamomile	
<b>General</b>	Family: Sunflower (Asteraceae) Introduced from Europe Colorado List B - Control required
<b>Common names</b>	Mayweed, scentless mayweed
<b>Habitat</b>	Forb Can live as an annual, biennial or short-lived perennial Found in meadows, pastures, rights-of-way, drainages
PLANT	
<b>Vegetation</b>	Plant to 3 feet tall Leaves ferny, odorless
<b>Roots</b>	Fibrous
<b>Flower</b>	June-October White with yellow center Centers tend to dome and the petals bend down as the flowers mature Single on each stem 3/4 - 1 inch wide
<b>Seed</b>	As many as 300,000 per plant May last up to 15 years in the soil
<b>Seedling</b>	Spring May form dense patches
<b>Reproduction</b>	Seed

<b>CONTROL</b>			
<b>BIOLOGICAL</b>	None		
<b>CHEMICAL</b>	<i>Stage</i>	<i>Herbicide</i>	<i>Timing</i>
	Rosette to bolting	Aminopyralid (Milestone) Chlorsulfuron (Telar) Metsulfuron (Escort XP)	Spring- early Summer
<b>CULTURAL</b>	Prevention Removal - make sure the entire root is removed. Must be repeated as new plants germinate Shallow tillage when plants are small, before flowers form		
<b>MECHANICAL</b>	<i>Stage</i>	<i>Method</i>	<i>Timing</i>
<b>Burning</b>	N/A		
<b>Grazing</b>	N/A	May cause blistering to livestock	
<b>Mowing</b>	N/A	Plants will regrow	

Use all chemicals according to the manufacturer's label.  
No specific recommendation or endorsement is made or implied by listing the above methods or products.

11/2015